

## **Review of the areal extent and the volume of the Serra Geral Formation, Paraná Basin, South America**

Heinrich Theodor FRANK, Márcia Elisa Boscato GOMES & Milton Luiz Laquintinie FORMOSO

Instituto de Geociências, Universidade Federal do Rio Grande do Sul, Caixa Postal 15.001, CEP 91501-970 Porto Alegre/RS, Brasil. E-mail: heinrich.frank@ufrgs.br; marcia.boscato@ufrgs.br; milton.formoso@ufrgs.br

Recebido em 04/2008. Aceito para publicação em 06/2009.  
Versão online publicada em 19/11/2009 ([www.pesquisasemgeociencias.ufrgs.br](http://www.pesquisasemgeociencias.ufrgs.br))

**Abstract** - In the discussion of the origin of continental flood basalt provinces, one of the most important constraints is the volume of the mobilized magma. Concerning the Serra Geral Formation in the Paraná Basin (South America), most authors cite an actual areal extent of 1,200,000 km<sup>2</sup> and a volume of approximately 800,000 km<sup>3</sup>. The same volume is often used for the Paraná-Etendeka Continental Flood Basalt Province, from which the Serra Geral Formation is the major part. We investigated the precision of these numbers through a complete historical review of their origin. We found that the areal extent cut off of 1,200,000 km<sup>2</sup> is only an estimate made in 1934, and that the volume cut off of 800,000 km<sup>3</sup> aroused using this areal extent estimate and an estimate of a mean thickness of 650 m of the flows made in 1966. Using a new map, we found that the area covered by the volcanic rocks in the Paraná Basin is of only 917,000 km<sup>2</sup> (+- 15,000 km<sup>2</sup>). With isopach maps of 1987 and 1990, we calculated the volume of the extrusive rocks to be of at least 450,000 km<sup>3</sup> and of the intrusive sill-type bodies to be of at least 112,000 km<sup>3</sup>. With these estimates, the volume of the Serra Geral Formation should be considered to be of more than 600,000 km<sup>3</sup>. For the Paraná-Etendeka Continental Flood Basalt Province as a whole, a volume of at least 1,700,000 km<sup>3</sup> should be considered.

**Keywords:** Serra Geral Formation, Paraná-Etendeka Continental Flood Basalt Province.

**Resumo** - REVISÃO DA EXTENSÃO AREAL E DO VOLUME DA FORMAÇÃO SERRA GERAL, BACIA DO PARANÁ, AMÉRICA DO SUL. A discussão sobre a origem das Províncias de Basaltos de Platô Continentais tem como um de seus elementos mais importantes o volume do magma mobilizado. Em relação às rochas da Formação Serra Geral na Bacia do Paraná (América do Sul), a maioria dos autores cita uma extensão de 1.200.000 km<sup>2</sup> e um volume de aproximadamente 800.000 km<sup>3</sup>. O mesmo volume é usado frequentemente para a Província de Basaltos de Platô Continental Paraná-Etendeka, da qual a Formação Serra Geral constitui a maior parte. Nós investigamos a precisão desses números através de uma revisão histórica completa acerca de sua origem. Verificamos que o valor relativo à extensão, de 1.200.000 km<sup>2</sup>, é apenas uma estimativa apresentada em 1934 e que o volume de 800.000 km<sup>3</sup> foi obtido multiplicando este valor de extensão por uma estimativa da espessura média dos derrames de lava de 650 metros apresentada em 1966. Usando um mapa recente, verificamos que a área coberta por rochas vulcânicas na Bacia do Paraná é de apenas 917.000 km<sup>2</sup> (+- 15.000 km<sup>2</sup>). Através de mapas de isópacos de 1987 e 1990, calculamos o volume das rochas extrusivas como sendo de pelo menos 450.000 km<sup>3</sup> e que o volume referente aos corpos intrusivos do tipo sill é de pelo menos 112.000 km<sup>3</sup>. Através destas estimativas, o volume de rochas da Formação Serra Geral deve ser considerado como sendo superior a 600.000 km<sup>3</sup>. A Província de Basaltos de Platô Continental Paraná-Etendeka como um todo possui um volume de no mínimo 1.700.000 km<sup>3</sup>.

**Palavras-chave:** Formação Serra Geral, Província de Basaltos de Platô Continental Paraná-Etendeka.

## 1. Introduction

Among the subaerial Large Igneous Provinces (LIPs) of the world, the Cretaceous Paraná-Etendeka Continental Flood Basalt Province (CFBP) ranks as second largest, surpassed only by the Siberian Traps in the Tunguska Basin. The origin of the Paraná-Etendeka CFBP is related to the opening of the southern Atlantic Ocean, but the constraints of the volcanic events are still controversial, as in other CFB Provinces too (Self *et al.*, 1996; Bondre *et al.*, 2004). The triggering mechanism and the heat source has been seen in the hot spot of Tristan da Cunha by some authors (e.g. Hill, 1991; Hawkesworth *et al.*, 1992; Wilson, 1993), but other contributions refuse the mantle plume influence in the volcanic event (Sheth, 1999; Ernesto *et al.*, 2002; Marques *et al.*, 2005). To refine the model, especially concerning alternative heat sources for the magma generation, one of the most important data concerns in the volume of magma produced during the volcanic activity. Despite hundreds of geochemical, geochronological, stratigraphical, petrographical, mineralogical and palaeomagnetic studies during the last decades, the areal extent and volume numbers presented for the Serra Geral Formation are highly variable. We made a complete historical review of the evolution of these numbers for the extrusive and intrusive rocks of the Serra Geral Formation in the Paraná Basin (South America) and calculated new values using available maps, now being able to define the precision of the used data and to present more realistic numbers.

## 2. Geological setting

The bulk of the Paraná-Etendeka CFBP is located in the Paraná Basin (Brazil, Argentina, Uruguay and Paraguay), an intracratonic basin covering 1,500,000 km<sup>2</sup>, developed between the Ordovician and the Cretaceous (Zalán *et al.*, 1987) (Fig. 1). The volcanic-sedimentary sequence is up to 7.5 km thick and is composed of six supersequences: Rio Ivaí, Paraná, Gondwana I, II and III and Baurú (Milani, 1997). The African part of the province is located in Etendeka-Namibia (Erlank *et al.*, 1984) and in Angola (Alberti *et al.*, 1992). A number of contributions outline the geochemical aspects of the volcanic rocks (e.g. Peate *et al.*, 1992), of which general aspects can

be found in Melfi *et al.* (1988). The volcanic rocks are called Serra Geral Formation in Brazil and Argentina, Arapey (flows) and Cuaró (sills) Formations in Uruguay, and Alto Paraná Formation in Paraguay. The most cited name, Serra Geral (White, 1908, v. 1, p. 17), derived from the name given to the eastern escarpment of the lava flows in the states of Paraná, São Paulo and Minas Gerais (Brazil). The age of the volcanic events was detailed through a number of contributions, becoming more accurate with the evolution of analytical methods. Turner *et al.* (1994) established an age of 137-127 Ma for the volcanic period.

## 3. Estimations for the Serra Geral Formation

### 3.1. Areal extent of the Serra Geral Formation

Two main figures are used to present the actual extension of the volcanic rocks of the Serra Geral Formation in the Paraná Basin. A value of 800,000 km<sup>2</sup> was established by Baker (1923) and referred to by much-cited contributions as Guimarães (1933), Sanford & Lange (1960), Almeida (1981) and Petri & Fúlfaro (1983). The number of 1,200,000 km<sup>2</sup> is an estimate made by Oppenheim (1934), who added to the volcanic outcroppings their probable extension in Uruguay and Argentina beneath post-Cretaceous sediments. This estimate was cited by Oliveira (1943) and Maack (1952), the latter being the most cited article with this number. Several other numbers, most of them of about 1,000,000 km<sup>2</sup>, can be found in a number of other contributions.

The original area covered by the Serra Geral flows is very difficult to establish and depends on the decision to interpret all basic dykes outcropping within basement rocks and in the sedimentary filling of the basin as feeders of lava flows, as assumed by Baker (1923). The original area was estimated to be 1,000,000 km<sup>2</sup> (Baker, 1923) or 2,000,000 km<sup>2</sup> (Renne *et al.* 1992). When defining the original perimeter of the lava flow field, isolated occurrences of basalts and diabases, resting on sedimentary rocks some tens of kilometers in front of the actual escarpment of the volcanic rocks, always have been seen as erosive remnants of lava flows and have been included. But most of them are exhumed sill-type bodies (Davino *et al.*, 1984), whose true nature was difficult to recognize due to its deep weathering and the scarcity of good

outcrops.

For the measuring of the actual area we defined a perimeter that includes the area of occurrence of the volcanic rocks and the overlying younger sediments. It is evident that a great and variable retreat of the border of the volcanic rocks has occurred. The error increased due to the very irregular limit approximately between Uberaba (state of Minas Gerais) and Botucatu (state of São Paulo), and due to the mapping of flows and sills as a single unit. Furthermore, the border of the outcropping volcanic rocks is covered by Cenozoic sediments of the Fray Bentos, Ituzaingó and Pampeana Formations in northeast Argentina, and by Cretaceous sedimentary rocks (Asencio, Mercedes, Guichón and Miguez Formations) in Uruguay.

The measured area includes (1) the area of outcropping basic volcanic rocks, (2) the area of outcropping acid volcanic rocks (64,000 km<sup>2</sup>, Nardy *et al.*, 2002) in the southeast corner of the basin, and (3) the areas inside of the perimeter of outcropping volcanics covered by post-volcanic sediments, like the Bauru Group (350,000 km<sup>2</sup>, Goldberg & Garcia, 2000, or 400,000 km<sup>2</sup>, Milani, 1997) and minor Formations: Cachoeirinha (Milani, 1997), Tupanciretã (Favilla *et al.*, 1995), Moinho (Böger *et al.*, 1993) and Missões (Consórcio Hidroservice-Hidrened, 1972), the last three in the state of Rio Grande do Sul, Brazil. The cartographic reference is the "Mapa de Integração Geológica da Bacia do Prata e Áreas Adjacentes" (Mercosul, 2001), on a scale of 1:2,500,000 and policonic projection. Considering the purposes of the review and the large number of uncertainties involved, a careful planimeter-based measurement was considered sufficient. A second apparatus was used for control, with each polygon measured at least four times, rendering a number of 917,000 km<sup>2</sup>, with an estimated error of 10,000-15,000 km<sup>2</sup>. In this way, considering an area of about 400,000 km<sup>2</sup> covered by post-volcanic sediments (Bauru Group and minor Formations), the remaining total area of exposed basalt is of approximately 500,000 km<sup>2</sup>, as estimated by Cabrera (1971, p. 23).

### 3.2. Volume of the Serra Geral Formation

The first volume calculations for the Serra Geral Formation were done by Baker (1923). These calculations become more precise through a number of contributions (e.g. Leinz, 1949; Teruggi, 1955; Leinz *et al.*, 1966). Volumes of

individual lava flows from the Paraná Basin are available only from Uruguay. One flow is cited by Bossi & Caggiano (1974) having a volume of >135 km<sup>3</sup>, and Bossi & Navarro (1991, v. 2, Tab. 14-6, p. 722) present a table of 38 flows ranging from 0.2 to 72 km<sup>3</sup>, with a total volume of at least 300 km<sup>3</sup>. The first "Isopach Map of the Basaltic Trapp in Paraná Basin", made by Leinz *et al.* (1966), does not represent the thickness of the lava flows, because it added to the thickness of the flows the thickness of the sills beneath them in the sedimentary rocks found during deep drillings (Bigarella & Salamuni, 1967, p. 30-31). Later, Leinz *et al.* (1968) presented a new map only with the thickness of the lava flows. The most cited value for the volume of the Serra Geral volcanics, of approximately 800,000 km<sup>3</sup>, was based in part on an estimate of a mean thickness of the lava flows of 650 m made by Leinz *et al.* (1966) and the areal extent estimate of 1,200,000 km<sup>2</sup>. However, this number did not take into account the intrusive bodies in the Paraná Basin and is absolutely not representative for the Province as a whole, since it did not consider the African part and the rocks that subsided in the marginal basins of South America and Africa.

The rock volumes lost horizontally and vertically by erosion process remain unknown. The lava flow pile of the Serra Geral Formation varies in thickness from few tens of meters at the borders of the basin to 1700 m (Almeida, 1986) near the city of Cuiabá Paulista (in the state of São Paulo), and scattered spots show thicknesses up to 2,5 km (Stanley *et al.*, 1985, fig. 10). The thickness of rocks removed by erosion was estimated, near the hydroelectric power plant of Itaipu, to be of 100-200 meters (Itaipu Binacional, 1994) and changes a lot in the basin due to the intense faulting and later erosion which leveled off the faulted blocks, as can be seen in the example of figure 2.

We obtained a reference value for the actual volume of the extrusive rocks of the Serra Geral Formation in the Brazilian part of the Paraná Basin through the "Isopach Map of the Extrusive Rocks of Serra Geral Formation" (Zalán *et al.*, 1987, Fig. 13), rendering at least 450,000 km<sup>3</sup> (Fig. 3). The number is of very low precision due to the scale of the map, and the occurrences in Uruguay, Argentina and Paraguay must be added. To underline the conservative character of the map, we brought a few new points indicated in figure 3: #a = 290 m (map indicates <100 m), #b = 1115 m (map indicates 400 m) and #c =

1000 m (map indicates 600 m). Data is from Machado (2005, #a, #b) and from B.L. Waichel (#c, personal comm.).

The intrusive sill-type bodies related to the Serra Geral Formation (Zalán *et al.*, 1985), emplaced in the sedimentary sequence of the Paraná Basin, show individual thicknesses of up to 400 meters and areas of up to 900 km<sup>2</sup> (Melfi *et al.*, 1988). The added thickness of the sills found through boreholes reach values of more than 1,000 meters and were compiled by Zalán *et al.* (1986) in the “Isopach Map of the Sill Type

Intrusions of Serra Geral Formation” (Fig. 4). The map refers only to the intrusives in the Brazilian part of the basin and can be refined with more data gathered during the last 20 years, still kept as internal reports of the oil companies. The evaluation of this map rendered us a minimum value of about 112,000 km<sup>3</sup> for these sill-type intrusives. In this way, we believe that the intrusive bodies in the basin as a whole represent volumes close to the entire Columbia River Basalt Group (174,300 km<sup>3</sup>, Tolan *et al.*, 1989).




Figure 1. Outline of the distribution of the Serra Geral rocks in the Paraná Basin, juxtaposed with the African counterparts in Namibia (Modified from Mercosul, 2001; African part after Hawkesworth *et al.*, 1992).




Figure 2. Geological section showing faulted blocks in Serra Geral volcanics in the states of Santa Catarina (SC) and Rio Grande do Sul (RS) (Mod. from Machado, 2005).




Figure 3. Isopach map of the extrusive rocks of the Serra Geral Formation (Mod. from Zalán *et al.*, 1986). “#” indicates locations of boreholes that provided some new data (see text).




Figure 4. Isopach map of sill-type bodies of the Serra Geral Formation (Mod. from Zalán *et al.*, 1986).

#### 4. Volume estimations for the Paraná-Etendeka CFBP

The volume of the Paraná-Etendeka Province as a whole is a sum of the volumes of the volcanics of the Paraná Basin and the related rocks in Etendeka (Namibia) and Angola, added to the ones now located in the continental margins of South America and Africa. A number of articles cite volumes for the Province between 800,000 km<sup>3</sup> (Peate *et al.*, 1990) and 2,350,000 km<sup>3</sup> (Gladzenko *et al.*, 1997).

In Africa, the rocks of the Etendeka Group in Huab Basin (Namibia) cover 78,000 km<sup>2</sup> (Erlank *et al.*, 1984) or 80,000 km<sup>2</sup> (Peate *et al.*, 1992, Hawkesworth *et al.*, 1992) with a maximum thickness of 900 meters, but the original thickness probably exceeded 2,000 meters (Reuning & Martin, 1957). Their volume, according to Milner *et al.* (1992), is of about 70,000 km<sup>3</sup>. In Angola (Alberti *et al.*, 1992), an unknown volume of rocks (Novo Redondo and

Lucira Formations) still awaits evaluation. The related volcanic rocks now situated in the marginal basins of Brazil and Africa (Fig. 5) are recognized as symmetrical provinces of seaward dipping reflectors (SDR), with the same composition, age and stratigraphical position as the Serra Geral Formation (Bueno, 2004). At the Brazilian margin, the volcanic rocks occur in the basins of Espírito Santo and Campos (Cabiúnas Formation), Santos (Camboriú Formation) and Pelotas (Imbituba Formation) (Fodor *et al.*, 1983; Fodor & Vetter, 1985; Almeida, 1986; Macedo, 1987; Macedo, 1989, p. 160 and fig. 5 to 7; Mizusaki *et al.*, 1992; Bueno, 2004), probably with volumes as large as the extrusive rocks in the Paraná Basin. At the African margin, in front of Namibia, the similar setting of the volcanic rocks is described by Gladzenko *et al.* (1997), who appoint volumes of 580,000 km<sup>3</sup> and 500,000 km<sup>3</sup> for the African and South American SDRs, respectively.




Figure 5. Tectonic evolution of Santos Basin (Brazilian platform) between the (A) Eocretaceous (130 Ma) and the (B) Neocretaceous (65 Ma) with the evolution of the extrusive rocks of the Paraná-Etendeka Continental Flood Basalt Province (in black) (Mod. from Macedo, 1987, fig. 10).

## 5. Discussion and conclusions

The discussion concerning the volumes of Serra Geral Formation and the Paraná-Etendeka CFBP is a complex task. After the volume estimate of Leinz *et al.* (1966, 1968) for the Serra Geral Formation, several hundred of deep drillings were made in the Paraná Basin, both in volcanic and sedimentary rock areas. In Brazil, the drillings are from PAULIPETRO (Consortium IPT-CESP), Companhia de Pesquisa de Recursos Minerais (CPRM - Geological Survey of Brazil) and PETROBRÁS (Petróleo Brasileiro SA). Some information from the drillings by PETROBRÁS was published by Northfleet *et al.* (1969) and by Zalán *et al.* (1990, p. 159, fig. 15). Other drillings were performed in Argentina by BRASPETRO (Petrobrás Internacional SA), in Paraguay by Texaco and PETROPAR (Petróleos Del Paraguay) and in Uruguay by ANCAP (Administración Nacional de Combustibles, Alcohol y Portland) and by the Instituto Geológico del Uruguay. These drillings, added to a wealth of offshore reports from Brazil, provided a great amount of new data, but almost all information is not available for public access.

The most cited number for the areal extent of the Serra Geral Formation (1,200,000 km<sup>2</sup>) is not consistent and should be avoided. Instead, we propose the value of 917,000 km<sup>2</sup> for the actual areal extent of the Serra Geral Formation in Paraná Basin, with some 500,000 km<sup>2</sup> of outcropping volcanic rocks. The estimated volume of the Serra Geral Formation rocks is of at least 600,000 km<sup>3</sup>. For the Paraná-Etendeka CFBP, a magma volume estimate of at least 1,700,000 km<sup>3</sup> is far more realistic.

With these new estimates, the much cited proportion of different rock types of the Serra Geral Formation (90% of tholeiitic basalts, 7% of tholeiitic andesites and 3% of rhyodacites-rhyolites, Melfi *et al.*, 1988) requires a revision, reducing the participation of the more acid rock types. Future works coupling the existent unavailable borehole and geophysical data of Paraná Basin will provide more accurate estimates of area and volume of the Serra Geral Formation.

**Acknowledgments** - We thank Norberto Dani, Heinrich Hasenack, Eliseu Weber, Jorge Luiz Barbosa da Silva, Carlos Augusto Sommer, Ubiratan Faccini and Nelson Rivaldo for their help. Thanks also to Breno

Leitão Waichel for helpful suggestions and to two anonymous reviewers.

## References

- Alberti, A., Piccirillo, E.M., Bellieni, G., Civetta, L., Comin-Chiaromonte, P. & Morais, E.A.A. 1992. Mesozoic acid volcanics from Southern Angola: petrology, Sr-Nd isotope characteristics and correlation with the acid stratoid volcanic suites of the Paraná basin (south-eastern Brazil). *European Journal of Mineralogy*, 4: 597-604.
- Almeida, F.F.M. 1981. Síntese sobre a Tectônica da Bacia do Paraná. In: Simpósio Regional de Geologia, 2., 1981, Curitiba, *Atas...*, Curitiba, SBG. v. 1, p.1-20.
- Almeida, F.F.M. 1986. Distribuição regional e relações tectônicas do magmatismo pós Paleozóico no Brasil. *Revista Brasileira de Geociências*, 16: 325-349.
- Baker, C.L. 1923. The lava field of the Paraná Basin, South America. *Journal of Geology*, 31: 66-79.
- Bigarella, A, J.J. & Salamuni, R. 1967. A Review of South American Gondwana Geology. In: IUGS (Eds.): *Reviews Prepared for the First Symposium on Gondwana Stratigraphy, Mar del Plata, Argentina*, p. 7-137.
- Böger, H. & Kowalczyk, G. 1993. Stratigraphische, Sedimentologische und Paläoökologische Untersuchungen im Mesozoikum der Depressão Periférica in Rio Grande do Sul, Brasilien. Geologisch-Paläontologisches Institut und Museum, Universität Kiel, *Berichte-Reports*, 63: 1-72.
- Bondre, N.R., Duraiswame, A. & Dole, G. 2004. Morphology and emplacement of flows from Deccan Volcanic Province, India. *Bulletin of Volcanology*, 66: 29-45.
- Bossi, J. & Caggiano, W. 1974. Contribuición a la geología de los yacimientos de amatistas en del departamento de Artigas (Uruguai). In: CONGRESSO BRASILEIRO DE GEOLOGIA, 28., 1974. Porto Alegre, *Anais...*, Porto Alegre, SBG, v. 3, p. 301-317.
- Bossi, J. & Navarro, R. 1991. *Geologia Del Uruguay*. Montevideo, Universidad de la Republica, 2 vol.
- Bueno, G.V. 2004. Diacronismo de eventos no rifte Sul-Atlântico. *Boletim de Geociências da Petrobrás*, 12(2): 203-229.
- Cabrera, J.G. 1971. *Geological and Engineering Properties of Basaltic Flows and Interbeds throughout the Upper Paraná Basin, Brazil*. 208p. PhD Thesis. Faculty of the Graduate School of Cornell University. Ann Arbor, Michigan.
- Consórcio Hidroservice - Hidrened. 1972. *Estudo do Aproveitamento do Trecho Limítrofe do Rio Uruguai e de seu Afluente Rio Pipiri-Guaçu, entre o Brasil e a Argentina*. Relatório de Inventário e Pré-Viabilidade (Etapas 1 e 2). Anexo 3 (Geologia).
- Davino, A., Sinelli, O. & Souza, A. 1984. Predominância de Formas Dômicas de Diabásio na Área da "Depressão Periférica" da Bacia do Paraná no

- Estado de São Paulo. In: CONGRESSO BRASILEIRO DE GEOLOGIA, 33., 1984. Rio de Janeiro. *Anais...*, São Paulo, SBG, v. 4, p. 2549-2562.
- Ernesto, M., Marques, L.S., Piccirillo, E.M., Molina, E.C., Ussami, N., Comin-Chiaramonti, P. & Bellieni, G. 2002. Paraná Magmatic Province - Tristan da Cunha plume system: fixed versus mobile plume, petrogenetic considerations and alternative heat sources. *Journal of Volcanology and Geothermal Research*, 118: 15-36.
- Erlank, A.J., Marsh J.S., Duncan, A.R., Miller R.McG., Hawkesworth, C.J., Betton, P.J. & Rex D.C. 1984. Geochemistry and petrogenesis of the Etendeka volcanic rocks from SWA/Namibia. *Geological Society of South Africa Special Publication*, 13: 195-245.
- Favilla, C.A.C., Rodrigues, T.L.N. & Caldasso, A.L.S. 1995. Depósitos Sedimentares Sin a Pós Vulcanismo Basáltico no Rio Grande do Sul. In: SIMPÓSIO SUL-BRASILEIRO DE GEOLOGIA, 6., Encontro de Geologia do Cone Sul, 1., 1995. Porto Alegre. *Anais...*, Porto Alegre, SBG, p. 267-269.
- Fodor, R.V. & Vetter, S.K. 1985. Mineral chemistry and petrography of passive-margin basalt Southeastern Brazil. *Revista Brasileira de Geociências*, 15(1): 36-47.
- Fodor, R.W., Mckee, E.H. & Asmus, H.E. 1983. K-Ar ages and the opening of the South Atlantic ocean: basaltic rocks from brazilian margin. *Marine Geology*, 54(1983/1984): M1-M8.
- Gladchenko, T.P., Hinz, K., Eldholm, O., Meyer, H., Neben, S. & Skogseid, J. 1997. South Atlantic volcanic margins. *Journal of the Geological Society*, 154: 465-470.
- Goldberg, K. & Garcia, A.J.V. 2000. Paleobiogeography of the Bauru Group, a dinosaur-bearing Cretaceous unit, northeastern Paraná Basin, Brazil. *Cretaceous Research*, 21: 241-254.
- Guimarães, D. 1933. Província Magmática do Brasil Meridional. *Boletim do Instituto Geológico e Mineralógico do Brasil*, 64: 1-70.
- Hawkesworth, C.J., Gallagher, K., Kelley, M., Mantovani, M.S., Peate, D.W., Regelous, M. & Rogers, N.W. 1992. Paraná magmatism and the opening of the South Atlantic. In: Storey, B.C.; Alabaster, T. & Pankhurst, R.J.(Eds). *Magmatism and the Causes of Continental Breakup*. London, Geological Society of London, p. 221-240 (Geological Society Special Publication, n. 68).
- Hill, R.I. 1991. Starting plumes and continental breakup. *Earth and Planetary Science Letters*, 104: 398-416.
- Itaipu Binacional. 1994. *Itaipu: Hydroelectric Project - Engineering Features*. Curitiba, PR.
- Leinz, V. 1949. Contribuição à geologia dos derrames basálticos do Sul do Brasil. *Boletim da Faculdade de Filosofia, Ciências e Letras da Universidade de São Paulo*, 103(5): 1-61.
- Leinz, V., Bartorelli, A., Sadowski, G.R. & Isotta, C.A.L. 1966. Sobre o comportamento espacial do trapp basáltico da Bacia do Paraná. *Boletim da Sociedade Brasileira de Geologia*, 15(4): 79-91.
- Leinz, V., Bartorelli, A. & Isotta, C.A.L. 1968. Contribuição ao Estudo do Magmatismo Basáltico Mesozóico da Bacia do Paraná. *Anais da Academia Brasileira de Ciências*, 40(Suplemento): 167-181.
- Maack, R. 1952. Die Entwicklung der Gondwana-Schichten Suedbrasiliens und ihre Beziehungen zur Karroo Formation Suedafrikas. In: INTERNACIONAL GEOLOGICAL CONGRESS OF ALGIERS, 19., 1952. Table de Matières, Algeries, IUGS, *Proceedings...*, v. 19, p.339-372 (Simposium sur les Séries de Gondwana).
- Macedo, J.M. 1987. Evolução Estrutural da Bacia de Santos e Áreas Continentais Adjacentes. In: SIMPÓSIO SUL-BRASILEIRO DE GEOLOGIA, 3., 1987, Curitiba (PR), *Anais...*, Curitiba, SBG, v. 2, p. 875-895.
- Macedo, J.M. 1989. Evolução Tectônica da Bacia de Santos e Áreas Continentais Adjacentes. *Boletim de Geociências da Petrobrás*, 3(3): 159-173.
- Machado, J.L.F., 2005. *Compartimentação e Arcabouço Hidroestratigráfico do Sistema Aquífero Guarani no Rio Grande do Sul*. São Leopoldo, 237p. Tese de Doutorado em Geociências, Universidade do Vale do Rio dos Sinos, São Leopoldo, Brasil.
- Mantovani, M.S.M., Stewart, K., Turner, S. & Hawkesworth, C.J. 1995. Duration of Paraná magmatism and implications for the evolution and sources regions of continental flood basalts. In: SYMPOSIUM ON THE PHYSICS AND THE CHEMISTRY OF THE UPPER MANTLE, 1995., São Paulo. *Proceedings...* Anais da Academia Brasileira de Ciências, 67(Suplemento 2): 163-170.
- Marques, L.S., Ernesto, M., Piccirillo, E.I.G., Figueiredo, A.M.G. & Min, A. 2005. Identificação de Diferentes Pulsos Magmáticos no Enxame de Diques Toleíticos da Serra do Mar. In: SIMPÓSIO DE VULCANISMO DE AMBIENTES ASSOCIADOS, 3., 2005, Cabo Frio, Rio de Janeiro, Brasil., *Anais...*, São Paulo, SBG. p. 289-293.
- Melfi, A.J., Piccirillo, E.M. & Nardy, A.J.R. 1988. Geological and magmatic aspects of the Paraná Basin (Brazil). An introduction. In: Piccirillo, E.M. & Melfi, A.J. (Eds.) *The Mesozoic flood volcanism of the Paraná basin: petrogenetic and geophysical aspects*. São Paulo, Instituto Astronômico e Geofísico, USP, P. 1-13.
- Mercosul 2001. Subgrupo de Trabalho 9. Comissão Temática de Geologia e Recursos Minerais. *Mapa de Integração Geológica da Bacia do Prata e Áreas Adjacentes*. Porto Alegre. CPRM, 1 mapa, 4 partes, col. Escala: 1:2.500.000.
- Milani, E.J. 1997. *Evolução Tectono-Estratigráfica da Bacia do Paraná e seu Relacionamento com a Geodinâmica Fanerozóica do Gondwana Sul-Occidental*. Porto Alegre. 2 v., Tese de Doutorado, Programa de Pós-graduação em Geociências, Instituto de Geociências, Universidade Federal do



- Rio Grande do Sul.
- Milner, S.C., Duncan, A.R. & Ewart, A. 1992. Quartz latite rheoignimbrite flows of the Etendeka Formation, north-western Namibia. *Bulletin of Volcanology*, 54: 200-219.
- Mizusaki, A.M.P., Petrini, R., Bellieni, G., Comin-Chiaramonti, P., Dias, J., De Min, A. & Piccirillo, E.M. 1992. Basalt magmatism along the passive continental margin of SE Brazil (Campos basin). *Contributions to Mineralogy and Petrology*, 111: 143-160.
- Nardy, A.J.R., Oliveira, M.A.F., Betancourt, R.H.S., Verdugo, D.R.H. & Machado, F.B. 2002. Geologia e Estratigrafia da Formação Serra Geral. *Geociências*, 21(1/2): 15-32.
- Northfleet, A.A., Medeiros, R.A. & Mühlmann, H. 1969. Reavaliação dos dados geológicos da Bacia do Paraná. *Boletim Técnico da Petrobrás*, 12 (3): 291-346.
- Oliveira, A.I. & Leonardos, O.H. 1943. *Geologia do Brasil*. Ministério da Agricultura, Rio de Janeiro. 813 p.
- Oppenheim, V. 1934. Rochas Gondwânicas e Geologia do Petróleo do Brasil Meridional. Departamento Nacional de Produção Mineral, *Boletim do Serviço de Fomento da Produção Mineral*, 5: 37-50.
- Peate, D.W., Hawkesworth, C.J., Mantovani, M.S. & Shukowsky, W. 1990. Mantle plumes and flood basalt stratigraphy in the Paraná Basin, South America. *Geology*, 18: 1223-1226.
- Peate, D.W., Hawkesworth, C.J. & Mantovani, S.M. 1992. Chemical stratigraphy of the Paraná lavas (South America): classification of magma types and their spatial distribution. *Bulletin of Volcanology*, 55: 119-139.
- Petri, S. & Fúlfaro, V.J. 1983. *Geologia do Brasil (Fanerozóico)*. São Paulo. T.A. Queiroz, Editora da Universidade de São Paulo, São Paulo, 631 p.
- Renne, P.R., Ernesto, M., Pacca, I.G., Coe, R.S., Glen, J.M., Prevot, M. & Perrin M. 1992. The age of the Paraná Flood Volcanism, rifting of Gondwanaland, and the Jurassic-Cretaceous boundary. *Science*, 258: 975-979.
- Reuning, E. & Martin, H. 1957. Die Prä-Karoo-Landschaft, die Karoo-Sedimente und die Karoo-Eruptivgesteine des südlichen Kaokofeldes in Südwestafrika. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie - Abhandlungen*, 91: 93-212.
- Sanford, R.M. & Lange, F.W. 1960. Basin-study approach to oil evaluation of Paraná Miogeosyncline, South Brazil. *Bulletin of the American Association of Petroleum Geologists*, 44: 1316-1370.
- Self, S., Thordarson, Th., Keszthelyi, L., Walker, G.P.L., Hon, K., Murphy, M.T., Long, P. & Finnemore, S. 1996. A new model for the emplacement of Columbia River basalts as large, inflated pahoehoe lava flow fields. *Geophysical Research Letters*, 23(19): 2689-2692.
- Sheth, H.C. 1999. A historical approach to continental flood basalt volcanism: insights into pre-volcanic rifting, sedimentation, and early alkaline magmatism. *Earth and Planetary Science Letters*, 168: 19-26.
- Stanley, W.D., Saad, A.R. & Ohofugi, W. 1985. Regional Magnetotelluric Surveys in Hydrocarbon Exploration, Paraná Basin, Brazil. *American Association of Petroleum Geologists Bulletin*, 69(3): 346-360.
- Teruggi, M. 1955. Los Basaltos Tholeiíticos de Misiones. Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Buenos Aires. Tomo XVIII, Notas do Museu. *Geologia* 70: 259-278.
- Tolan, T.L., Reidel, S.P., Beeson, M.H., Anderson, J.L., Fecht, K.R. & Swanson, D.A. 1989. Revisions to the estimates of the areal extent and volume of the Columbia River Basalt Group. In: Reidel, S.P. & Hooper, P.R. (Eds.), *Volcanism and Tectonism in the Columbia River Flood-Basalt Province*. Boulder, Colorado. Geological Society of America, 1989. p. 1-20 (Geological Society of America, Special Paper 239).
- Turner, S., Regelous, M., Kelley, S., Hawkesworth, C.J. & Mantovani, M.S.M. 1994. Magmatism and continental break-up in the South Atlantic: high precision  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  geochronology. *Earth and Planetary Science Letters*, 121: 333-348.
- White, I.C. 1908. Relatório sobre as "coal measures" e rochas associadas do Sul do Brasil. *Relatório Final da Comissão de estudos das minas de carvão de pedra do Brasil*, Rio de Janeiro, Imprensa Nacional, Rio de Janeiro, 28., 617 f.
- Wilson, M. 1993. Magmatism and the geodynamics of basin formation. *Sedimentary Geology*, 86: 5-29.
- Zalán, P.V., Conceição, J.C., Astolfi, M.A.M., Appi, V.T., Wolff, S., Vieira, I.S. & Marques, A. 1985. Estilos Estruturais Relacionados a Intrusões Magmáticas Básicas em Rochas Sedimentares. *Boletim Técnico da Petrobrás*, 28(4): 221-230.
- Zalán, P.V., Conceição, J.C.J., Wolff, S., Astolfi, M.A.M., Vieira, I.S., Appi, V.T., Neto, E.V.S., Cerqueira, J.R., Zanutto, O.A., Paumer, M.L. & Marques, A. 1986. *Análise da Bacia do Paraná*. Relatório Interno da Petrobrás. Depex/Cenpes nº 1035-5765, Rio de Janeiro, 5 v.
- Zalán, P.V., Wolff, S., Conceição, J.C.J., Astolfi, M.A.M., Vieira, I.S., Appi, V.T. & Zanutto, O.A. 1987. Tectônica e Sedimentação da Bacia do Paraná. In: SIMPÓSIO SUL BRASILEIRO DE GEOLOGIA, 3., 1987. Curitiba, Atas..., Curitiba, SBG, 1987. v. 1, p. 441-477.
- Zalán, P.V., Wolff, S., Conceição, J.C.J., Marques, A., Astolfi, M.A.M., Vieira, I.S., Appi, V.T. & Zanutto, O.A. 1990. *Bacia do Paraná. Origem e Evolução de Bacias Sedimentares*. PETROBRÁS/SEREC/CEN-SUD, Rio de Janeiro, p. 135-168.